



Anti-parasitic effects of plant secondary metabolites on swine nematodes

Williams, A.R.; Pena-Espinoza, Miguel Angel; Fryganas, Christos; Ropiak, H.M.; Ramsay, Aina; Mueller-Harvey, I.; Thamsborg, S. M.

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Williams, A. R., Pena-Espinoza, M. A., Fryganas, C., Ropiak, H. M., Ramsay, A., Mueller-Harvey, I., & Thamsborg, S. M. (2014). *Anti-parasitic effects of plant secondary metabolites on swine nematodes*. Abstract from 13th International congress of parasitology, Mexico.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Anti-parasitic effects of plant secondary metabolites on swine nematodes

*Williams, Andrew*¹, *Peña, Miguel*², *Fryganas, Christos*³, *Ropiak, Honorata*³, *Ramsay, Aina*³, *Mueller-Harvey, Irene*³, *Thamsborg, Stig*¹

¹Department of Veterinary Disease Biology, University of Copenhagen, Denmark, ² National Veterinary Institute, Technical University of Denmark, ³SAPD, Reading University, UK.

BACKGROUND: Organic animal production presents challenges to animal health and productivity. In organic pig production, animals must have access to outdoor pastures which increases exposure to gastrointestinal parasites. Moreover, the routine use of synthetic anti-parasitic drugs is not allowed. Thus, novel parasite-control options are required. We present results from a comprehensive *in vitro* screen of plant secondary metabolites (PSM) from diverse plant sources on the economically important pig parasites *Ascaris suum* and *Oesophagostomum dentatum*.

METHODS: We focused on two PSM classes commonly found in natural diets – condensed tannins (CT) and sesquiterpene lactones (SL). Different CT-types were purified from various plant sources to reflect their diversity; SL were purified from forage chicory. These PSM were tested in inhibition assays of worm motility and migratory ability.

RESULTS: CT had potent activity against *A. suum*, with substantial inhibition of migration of *in vitro* hatched larvae (EC₅₀ values ranging from 40 to 120 µg/mL). In contrast, migratory ability of *O. dentatum* larvae was not significantly affected. However, the motility of adult *O. dentatum* was reduced after *in vitro* incubation with CT. The purified chicory extract showed potent inhibition of *A. suum* larval migration (EC₅₀ value of 42 µg/mL) and was also active against larval and adult stages of *O. dentatum*. Electron microscopy demonstrated significant structural damage to the cuticle and digestive tissues in nematodes exposed to PSM.

CONCLUSIONS: Plants rich in PSM such as CT and SL show promise as natural anthelmintics against two highly prevalent swine parasites. Experiments to determine *in vivo* efficacy and mechanisms of nematocidal action are on-going.